

2017

Injuries caused by pets in Asian urban households: A cross-sectional telephone survey

Emily Y. Y. Chan

The Chinese University of Hong Kong, emily.chan@cuhk.edu.hk

Yang Gao

Hong Kong Baptist University, gaoyang@hkbu.edu.hk

Liping Li

Shantou University

Po Yi Lee

The Chinese University of Hong Kong

This document is the authors' final version of the published article.

Link to published article: <http://dx.doi.org/10.1136/bmjopen-2016-012813>

Recommended Citation

Chan, Emily Y. Y., Yang Gao, Liping Li, and Po Yi Lee. "Injuries caused by pets in Asian urban households: A cross-sectional telephone survey." *BMJ Open* 7.1 (2017): e012813.

BMJ Open Injuries caused by pets in Asian urban households: a cross-sectional telephone survey

Emily Y Y Chan,¹ Yang Gao,^{2#} Liping Li,³ Po Yi Lee¹

To cite: Chan EYY, Gao Y, Li L, *et al.* Injuries caused by pets in Asian urban households: a cross-sectional telephone survey. *BMJ Open* 2017;7:e012813. doi:10.1136/bmjopen-2016-012813

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2016-012813>).

YG and EYYC contributed equally.

Received 30 May 2016
Revised 2 November 2016
Accepted 10 November 2016



CrossMark

For numbered affiliations see end of article.

Correspondence to
Professor Emily YY Chan;
emily.chan@cuhk.edu.hk

ABSTRACT

Objectives: Little is known about pet-related injuries in Asian populations. This study primarily aimed to investigate the incidence rate of pet-related household injuries in Hong Kong, an urban Chinese setting.

Setting: Cantonese-speaking non-institutionalised population of all ages in Hong Kong accessible by telephone land-line.

Participants: A total of 43 542 telephone numbers were dialled and 6570 residents successfully completed the interviews.

Primary and secondary outcome measures: Data of pet-related household injuries in the previous 12 months, pet ownership and socio-demographic characteristics were collected with a questionnaire. Direct standardisation of the incidence rates of pet-related household injuries by gender and age to the 2009 Hong Kong Population Census was estimated. Univariate and multivariate analyses were performed to estimate risks of socio-demographic factors and pet ownership for the injury.

Results: A total of 84 participants experienced pet-related household injuries in the past 12 months, with an overall person-based incidence rate of 1.28%. The majority of the victims were injured once (69.6%). Cats (51.6%) were the most common pets involved. Pet owners were at an extremely higher risk after controlling for other factors (adjusted OR: 52.0, 95% CI 22.1 to 98.7). Females, the unmarried, those with higher monthly household income and those living in lower-density housing were more likely to be injured by pets.

Conclusions: We project a pet-related household injury incidence rate of 1.24% in the general Hong Kong population, with 86 334 residents sustaining pet-related injuries every year. Pet ownership puts people at extremely high risk, especially the unmarried. Further studies should focus on educating pet owners to reduce pet-related injuries in urban Greater China.

BACKGROUND

Human–animal contact is a daily occurrence for most pet owners globally. In urban settings, dogs and cats are the most commonly involved animals.^{1–4} Urban pet ownership is a rising global trend along with economic

Strengths and limitations of this study

- This study may be the first of its kind focusing on pet-related injuries in Asian urban households.
- Though pet-related household injury incidence rate was as low as 1.24%, the large sample size (n=6570) supported reliable results.
- The cross-sectional design cannot draw a conclusion on any cause–effect relationship.
- The self-reported pet-related injuries may be subject to potential recall bias.
- Limited collection about the details of pet ownership in this study is constrained to develop effective protection measures.

development. China follows this urban trend for dog and cat ownership, and has the third largest dog population and the sixth largest cat population in the world.^{5 6} The latest available data have shown that China's pet population has grown by 20% in 5 years from 1999 to 2004, largely attributable to the rapid increase in urban areas.⁶ Given its rapid economic development in the past decades, especially the increased size of the population that has more disposable income resulting in affordable pet ownership, pet-related injuries would be expected to increase along with the rising pet ownership rate in urban China.

In the urban settings of high income countries, although victims with moderate to severe pet-related injuries may seek medical attention either in an emergency department or in a physician's office, most minor injuries go unreported, with victims administering first aid by themselves and then often do not seek or require medical attention.^{3 7} In Western high-income communities, young children aged 5–9 years bear most reported pet-related injuries, and dogs are the most commonly involved animals.^{1–3 8 9} Possession of a pet is found to be associated with a higher risk for animal bites. Adults with two or more dogs in the household were five



times more likely to be bitten than those living without dogs at home.⁴ In addition, pets are also associated with non-fatal fall-related injuries. An estimated average of 86 629 fall injuries each year were associated with cats and dogs, for an average annual injury rate of 29.7 per 100 000 population in the USA.⁴ Although there are various case reports and clinic-setting based studies, no studies of pet-related injuries have been conducted among the general population in Asia. There is a knowledge gap in the epidemic of pet-related injuries in this population. Without the knowledge, it is difficult to develop effective preventive strategies.

Hong Kong, located in South China, is a typical well-developed urban setting in Asia. Its social and economic developments as well as people's lifestyle have been referred or even copied by many other Asian cities. The pet ownership in Hong Kong is about 20%, relatively lower than in the Western developed countries (eg, 34.6% in Australia and 36.5% in the USA), mainly due to the limited living space and hurried lifestyle in Hong Kong.⁷ The knowledge of pet-related injuries in Hong Kong is scarce. A 2008 survey projected that annually 5400 animal bite-related injuries, which were serious enough to limit the normal activities of a person, occurred in the entire Hong Kong population.¹⁰ That study did not differentiate pets and non-pet animals. In addition, no further information such as incidence rate, distribution in the population and risk factors was reported. This study primarily aimed to investigate the incidence rate of non-fatal pet-related household injuries in Hong Kong. In addition, we also examined the associations of the socio-demographic characteristics and pet ownership with the injury. Results of the study provided better understanding about pet-related injuries in urban Asia, which can be used to develop effective strategies to reduce pet-related injuries in Asian population.

METHODS

Participants

A cross-sectional, population-based telephone survey was conducted during April to June, 2009. The target population was the Cantonese-speaking non-institutionalised population of all ages in Hong Kong. The Hong Kong population in 2009 was 6.97 million, of which about 95.8% were Cantonese speakers or able to use Cantonese.¹¹ Over 95% of the households in Hong Kong had a land-line telephone.^{12 13} Telephone numbers were randomly selected from current telephone directories. Telephone calls were made during weekday evenings (6:30 pm to 10:00 pm, Monday to Friday) and during daytime on the weekends (Saturday and Sunday) in order to avoid an over-representation of people not working. An eligible member (defined as a Chinese-speaking Hong Kong resident without hearing impairment) was selected from each of the contacted households. If more than one member was eligible in the same household, the one whose birthday closest to

the survey date was invited to participate into the study. If the selected member was under 16 years old, the adult member who knew the selected person most was invited to respond to the telephone interview. At least five calls were dialled at different time slots before we considered the number invalid and gave up. All telephone interviews were conducted by trained investigators. Verbal consent was obtained from the adult participants and parental verbal consent was sought for those aged under 18.

Sample size estimation

The telephone survey had a focus on all types of injuries occurring in household settings. Thus, the sample size was estimated based on the incidence rate of all non-fatal unintentional household injuries. Our pilot study, conducted in May–June 2007 among 1001 participants, indicated an incidence rate of all household injuries of 25%. After calculation, a sample size of 6500 yielded a 95% CI with a precision of 1.0% for an estimated rate of 25% ($\beta=0.80$ and $\alpha=0.05$).

Data collection and measures

A modified Chinese-language questionnaire based on the WHO's Guidelines for Conducting Community Surveys on Injuries and Violence was used.¹⁴ The questionnaire has been proved to be reliable and valid in our previous studies.^{12 15} All participants were asked about the following items.

Pet-related household injury: it was defined as unintentional, non-fatal physical damage to the human body caused by pets (eg, pet bites, pet scratches, falls caused by pets, etc), which took place in a household environment, including in the common areas of their residence such as clubhouses, common entry ways, car parks, stairwells or inside an elevator. A pet in this study was defined as any domesticated or tamed animal that is kept as a companion and cared for affectionately, including cats, dogs, rabbits, birds, guinea pigs, hamsters and so forth. Types and frequencies of this type of injury in the previous 12 months were asked. For those who indicated a pet-related injury as the most recent injury, the following information was further collected from them: day and place of that injury, body part(s) injured, human–pet interaction before the injury, ownership of the pet involved (pet ownership was defined as the pet owned by the family of the injured person), vaccination of that pet (eg, rabies vaccination, vaccination against distemper, parvovirus, hepatitis, leptospirosis and kennel cough for dogs and vaccination against feline enteritis for cats), whether attacked by the same pet before, treatment of that injury, absence from work or school caused by the injury and recovery time of the injury.

Socio-demographic and other factors: gender, age, marital status, education attainment, occupation and employment status, monthly household income, family size, type and size of the living area (square feet, ft²), living

density (ft²/person) and time spent at home (hours/day) were collected from the participants.

Statistical analysis


Data were analysed with the IBM SPSS Statistics (V.20). Overall, gender-specific and age-specific incidence rates of pet-related injuries in the previous 12 months were calculated. Direct standardisation of the incidence rate by gender and age was used to project the incidence of pet-related injuries in the entire Hong Kong population, indicated by the number of people injured and the overall incidence rate.¹⁶ χ^2 test was performed to analyse between-group differences in pet-related injuries. Univariate and multivariate logistic regression were fitted to estimate the risks of the socio-demographics and other confounding factors for the injury before and after adjustment for each other. In the univariate analyses, Bonferroni correction was applied to counteract the problem of multiple comparisons. Significance levels of $\alpha=0.0125$ (0.05/4) and $\alpha=0.00417$ (0.05/12) were therefore adopted respectively in comparisons between the study participants and the Hong Kong population (2009) and those between injured and non-injured participants. In the multivariate analyses, a hierarchical approach was adopted, with first regressing the socio-demographic factors (Model 1), and then entering pet ownership (Model 2). All socio-demographic variables in Model 1 were forwardly selected (Wald method) using $p<0.10$ as the entry and removal criteria. OR and 95% CI were then derived.

RESULTS

A total of 43 542 telephone numbers were dialled, in which 12 479 were successfully reached and 10 380 were with people eligible for the study (figure 1). Among the eligible people, 6723 (64.8%) agreed to participate into the study and were interviewed and 6570 (63.3%) successfully completed the interviews and with their data analysed. It was found a minor but significant difference (using $\alpha=0.0125$) in age distribution between the study participants and the general Hong Kong population, with our participants having slightly more children and young adults (table 1). In addition, our study had significantly more participants with higher education levels and doing a white-collar job, compared with the general population.

The overall person-based incidence rate of pet-related household injuries was 1.28% (n=84). Seventy-nine injured participants reported the injury frequency in the past 12 months, including 69.6% (n=55) experienced such injuries once, 17.7% (n=14) did twice or three times and 12.7% (n=10) did four times or more. The remaining five injured persons were unable to remember the number of injuries. Only one injury was caused by a related fall. The direct age/gender standardisation of rates estimated that annually 86 334 Hong Kong residents experienced pet-related household injuries, with an overall incidence rate of 1.24%.

There were significant differences in the distribution of age, marital status, size of living area and pet ownership according to the significance level of 0.00417 (table 2). Of the injured participants, 64 (76.2%)



*: ineligible respondents included non-Chinese-speaking persons, non-Hong Kong residents, and hearing impaired persons (unable to be interviewed by phone calls).

#: incomplete interviews.

Figure 1 Recruitment process of study participants.

Table 1 Comparisons of major socio-demographic characteristics between the study participants and the Hong Kong population 2009

	Study participants		Hong Kong population (2009)†		p Value of χ^2 test
	n	%	n	%	
Gender (N=6561)					0.042
Male	3173	48.4	3 284 800	47.7	
Female	3388	51.6	3 688 000	52.3	
Age (years, N=6503)					<0.001
0–14	887	13.6*	863 800	12.4*	
15–24	906	13.9*	885 400	12.7*	
25–44	2070	31.8*	2 236 90	32.1*	
45–64	1837	28.2*	2 088 100	29.9*	
≥65	803	12.3*	898 600	12.9*	
Education in population aged 15 or above(N=5559)					<0.001
Primary or below	932	16.8*	1 589 800	22.8*	
Secondary	2,994	53.9*	3 611 900	51.8*	
Postsecondary‡	1633	29.4*	1 771 100	25.4*	
Occupation in working population (N=2873)					<0.001
White-collar job§	2376	82.7*	2 351 200	67.7*	
Blue-collar job¶	462	16.1*	1 111 500	32.0*	
Others	35	1.2*	5000	0.1*	

*Significant difference reached in the χ^2 test by using the actual number of population. Bonferroni correction was applied to correct multiple testing. A significance level of $\alpha=0.0125$ (0.05/4) was therefore adopted.

†Data sources: Hong Kong annual digest of statistics 2012 edition.

‡Postsecondary education included non-degree and degree courses.

§White-collar job: including manager and administrators, professionals, associate professionals, clerical support workers and service and sales workers.

¶Blue-collar job: including craft and related workers, plant and machine operators, elementary occupations and skilled agricultural and fishery workers.

reported a pet-related injury as the most recent injury. No significant difference in the distribution of any factor was found between the two groups (those who were injured by a pet and those who reported a pet-related injury as the most recent injury). **Table 3** provides the results of the pet-related injuries among those reporting a pet-related injury as their most recent injury. Hands (79.7%) were the most common body part injured. Cats (51.6%) and dogs (42.2%) were the most common pets involved. The majority of the involved pets belonged to the victims (92.2%) and had ever received vaccinations beforehand (70.3%). The majority of the injured participants (62.5%) dealt with the injury by themselves. Only one case sought medical care at a Chinese medicine clinic, with a total expenditure of HK \$330 (around US\$42.5) for two visits, consisting of HK \$310 (around US\$40.0) for the medical cost and HK\$20 (around US\$2.5) for the transport cost.

Table 4 presents incidence rates and risk estimates of socio-demographic factors and pet ownership for the injury. In the univariate analyses, young adults (15–24 years old, with age \geq 45 as reference), the unmarried, those with the highest monthly household income (\geq HK\$30 000), those living in the biggest house (>700 ft²) and pet owners were significantly more likely to be injured (using $\alpha=0.00417$), while retired participants were significantly less likely to experience

pet-related injury (with those in a white-collar job as reference, $\alpha=0.00417$). In the multivariate Model 1, where the selected socio-demographic factors were controlled for each other, females, the unmarried, those with higher monthly household income and those with lower density residences were significantly at higher risks for the injury ($\alpha=0.05$ for all multivariate analyses). In addition, increasing risks of monthly household income and living density were observed for pet-related injury. After controlling for pet ownership (Model 2), only the elevated risks of unmarried participants and the highest monthly household income (\geq HK\$30 000) remain significant (adjusted OR=1.80, 95% CI 1.04 to 3.10, $p=0.035$ and adjusted OR=7.60, 95% CI 1.01 to 57.1, $p=0.049$ respectively). In addition, pet owners were at extremely higher risk for the injury (adjusted OR=52.0, 95% CI 22.1 to 122.4, $p<0.001$).

Considering that it is a strong factor, we further performed univariate and multivariate logistic regression after stratification by pet ownership. Among non-pet owners, no socio-demographic characteristic was significantly related to pet-related injuries in χ^2 and multivariate analyses. Among pet owners, univariate logistic regression revealed that age group, marital status and living density were significantly related to pet-related injuries, while only marital status remains significance in the multivariate model, with unmarried participants

Table 2 Characteristics of the participants by pet-related injury in the past year

	No pet-related injury (N=6481)		Pet-related injury (N=84)		Pet-related injury as the latest injury (N=64)	
	n	%	n	%	n	%
Gender						
Male	3140	48.5	33	39.3	29	45.3
Female	3337	51.5	51	60.7	35	54.7
Age (years)						
0–4	198	3.1*	4	4.9*	1	1.6
5–14	676	10.5*	9	11.0*	6	9.4
15–24	884	13.8*	22	26.8*	17	26.6
25–44	2042	31.8*	28	34.1*	26	40.6
≥45	2621	40.8*	19	23.2*	14	21.9
45–64	1818	28.3*	19	23.2*	14	21.9
≥65	803	12.5*	0	0.0*	0	0.0
Marital status						
Married/cohabiting	3489	54.3*	30	36.6*	22	34.9
Unmarried (single/divorced/widowed)	2936	45.7*	52	63.4*	41	65.1
Education						
Primary or below	1447	23.0	9	11.0	6	9.5
Secondary	3206	51.0	46	56.1	31	49.2
Postsecondary†	1628	25.9	27	32.9	26	41.3
Occupational status						
White-collar job‡	2370	37.2	41	49.4	34	54.0
Blue-collar job§	469	7.4	2	2.4	1	1.6
Housewives	846	13.3	8	9.6	5	7.9
Retired	921	14.5	3	3.6	3	4.8
Students	1444	22.7	25	30.1	17	27.0
Others	315	4.9	4	4.8	3	4.8
Monthly household income (HK\$¶)						
<10 000	942	18.6	3	4.5	4	7.7
10 000–19 999	1422	28.0	19	28.4	16	30.8
20 000–29 999	1035	20.4	12	17.9	4	7.7
≥30 000	1677	33.0	33	49.3	28	53.8
Family size (persons)						
2–4	4732	73.7	62	75.6	48	75.0
≥5	1325	20.6	17	20.7	13	20.3
Living along (1 person)	364	5.7	3	3.7	3	4.7
Type of housing						
Public housing	1826	28.3	16	19.0	11	17.2
Subsidised housing	993	15.4	12	14.3	9	14.1
Private housing	3555	55.1	54	64.3	42	65.6
Others	77	1.2	2	2.4	2	3.1
Size of living area (ft²§)						
≤400	1645	27.7*	13	18.3*	11	20.0
401–600	1932	32.5*	14	19.7*	15	27.3
601–700	901	15.2*	16	22.5*	9	16.4
>700	1463	24.6*	28	39.4*	20	36.4
Living density (ft²/person§)						
<120	1511	25.5	9	12.9	9	16.4
120–159	1288	21.8	13	18.6	11	20.0
160–249	1985	33.5	29	41.4	21	38.2
≥250	1136	19.2	19	27.1	14	25.5
Time spent at home (hours/day)						
≤12	3074	47.8	40	48.2	32	50.0
13–18	2629	40.8	34	41.0	23	35.9
>18	734	11.4	9	10.8	9	14.1

Continued

Table 2 Continued

	No pet-related injury (N=6481) n %		Pet-related injury (N=84) n %		Pet-related injury as the latest injury (N=64) n %	
Pet ownership						
No	5675	87.9*	11	13.1*	5	7.8
Yes	780	12.1*	73	86.9*	59	92.2

*Significant difference level at $p=0.05$ reached in the χ^2 test. Bonferroni correction was applied to correct multiple testing. A significance level of $\alpha=0.00417$ ($0.05/12$) was therefore adopted. No significant differences were found between pet-related injury and pet-related injury as the latest injury.

†Postsecondary education included non-degree and degree courses.

‡White-collar job: including manager and administrators, professionals, associate professionals, clerical support workers and service and sales workers.

§Blue-collar job: including craft and related workers, plant and machine operators, elementary occupations and skilled agricultural and fishery workers.

¶Annual exchange rate in 2009: HK\$1=US\$0.129 (US\$1=HK\$7.75); HK\$10 000=US\$1290; HK\$20 000=US\$2580; HK\$30 000=US\$3870.

§1 ft²=0.093 m² (1 m²=10.76 ft²).

being at higher risk (adjusted OR=2.25, 95% CI 1.28 to 3.96, $p=0.005$).

DISCUSSION

This study found a person-based incidence rate of pet-related household injury of 1.28% in the previous 12 months among 6570 residents in Hong Kong. Using the direct age/gender standardisation rates, an overall incidence rate of 1.24% was projected for the entire Hong Kong population, with an estimation of 86 334 residents injured by pets annually. In our participants, the majority of the injured participants were bitten by pets once (69.6%), with hands as the body part most commonly injured (79.7%), and 98.4% did not seek medical attention. Cats (51.6%) were the most common pets causing injury, followed by dogs (42.2%). Most of the pets belonged to the injured (92.2%). Pet owners were at extremely higher risk for such injuries. In addition, females, the unmarried, those with higher monthly household income and those living in lower density housing were more likely to be injured by pets.

This study may be the first of its kind focusing on pet-related household injuries in the general population in an urban Chinese setting. We estimated that annually 1.24% of Hong Kong residents sustained pet-related household injury, which is similar to those reported by other developed countries.^{7 17} However, the projected size of people injured by pets (86 334) is much higher than the estimate from a previous study in Hong Kong on all causes of unintentional injuries ($n=5400$).¹⁰ Such differences may be mainly due to the different scopes of an 'injury', as defined between the two studies: we investigated all pet-related injuries while the previous study only focused on injuries serious enough to limit the normal activities of a person. Most of the cases in our study did not seek any medical attention or did not take sick leave, reflecting that most pet-related injuries in Hong Kong are minor, therefore, were not captured by the previous study. One limitation of the study is

over-representation of residents with higher education levels and doing white-collar jobs. One plausible explanation is due to the ability to make a connection during the times called. In Hong Kong, less educated people or those doing blue-collar jobs may work longer time or in the evening, which may have reduced their chances to be reached in this study. As education and occupation were not significantly related to pet-induced injury (table 4), the over-representation may not have much influence on the projected size of injured residents.

Consistent with many previous studies, we found pet ownership to be a strong risk factor for pet-related injuries.^{3 4 7 9 18} However, the risk estimation of pet ownership in this study (adjusted OR=52.0) is much higher than the ones reported by these previous studies, which may be attributable to the discrepancies between studies, such as place of the injuries (the household setting in this study vs all settings in other studies), severity of a pet-related injury interested (eg, injuries received medical care in most previous studies), types of pets interested (eg, dogs only in most previous studies) and so on. Nevertheless, the increased number of pet owners, along with the fast-paced economic development in China, could result in increase of pet-related injuries.^{5 6} Further studies should be conducted to confirm our findings and develop effective interventions to protect people from pet-related injuries.

In this study, females were more likely to be injured by a pet, which was consistent with the finding from the previous Hong Kong study,¹⁰ but contradictory with the results from studies in other countries.^{3 7 18 19} It is not clear about the cause of the gender difference among Hong Kong residents and other populations. One possibility is that the two studies in Hong Kong reported on any pet-related injuries while the other studies focused only on dog-induced injuries. Further analysis in this study indicated that males were slightly more likely to have dog-induced injuries (51.9% vs 48.1%), which may at least partly explain the difference. In contrast to the results from other studies, which showed that children

Table 3 Further information of the participants with pet-related injury as the latest injury (N=64)

	N	Per cent
Occurrence place of the injury		
Living room	47	73.4
Bedroom	7	10.8
Bathroom	5	7.8
Others	5	7.8
Occurrence day of the injury		
Weekday	30	46.9
Weekend and holiday	23	35.9
Don't know	11	17.2
Body part injured‡		
Hands	51	79.7
Legs	10	15.6
Arms	3	4.7
Head	2	3.1
Body	1	1.6
The pet involved		
Cat	33	51.6
Dog	27	42.2
Others†	4	6.2
The pet ever received any vaccination		
No	11	17.2
Yes	45	70.3
Don't know	8	12.5
Owner of the pet related		
Yes	59	92.2
No	5	7.8
Interaction before attack		
Playing with the pet	38	59.4
Cleaning the pet	6	9.4
Training the pet	5	7.8
Others	15	23.4
Ever attacked by the same pet before		
Yes	29	45.3
No	35	54.7
Treatment of the injury		
None	15	23.4
Self-treatment	40	62.5
Others (including medical care and treated by other persons)	9	14.1
Absence from work or school caused by the injury		
No	13	20.3
Yes	51	79.7
<1 day	48	94.1
≥1 day	3	5.9
Recovery time of the injury (days)		
<3	45	70.3
3–7	13	20.3
>7	6	9.4

‡As there were three injuries in which two body parts of the injured participants were injured, the sum of "n" is 67, not 64. However, the denominator is still the number of injured persons (N=64), so the sum of "%" is more than 100%.

†Others included rabbits, birds, guinea pigs, hamsters and so forth.

(especially those aged 5–9 years) were the most vulnerable cohort to sustain a pet-related injury,^{3 7 10 18 19} we found that young adults (aged 15–24 years) were at the highest risk among all the age cohorts. These

differences could be explained by different data sources used across the studies. The results from the other studies were based on data collected in clinics and emergency departments. Adults are less likely than children to seek medical care in clinical settings after being injured by pets, especially when the injuries are minor, which might have resulted in an underestimation of pet-related injuries among adults in those studies (due to selection bias). However, data in our study were collected from a representative sample of the general population, which may have avoided such selection bias, therefore, reflects the real situation of pet-related injuries more accurately.

We found that higher monthly household income, greater housing living space and larger living area per capita (lower living density) were risk factors for pet-related injuries, while those having blue-collar jobs were less likely to be injured. These associations suggest affluence as a risk factor, which is consistent with other studies.⁷ When controlling for each other, the associations of occupation status and housing size with an injury were no longer significant, while monthly household income and living density remained significant and were positively related with the injuries. This may suggest that income and living density may be better indicators of affluence in urban Asian settings. In addition, we also found that unmarried participants were twice as likely to sustain a pet-related injury compared to married participants and the relationship did not change much after controlling for other factors. The reason is unknown as no previous study reported significant relationship between marital status and pet-related injuries. The most common reason for pet ownership is to get a companion to alleviate loneliness.⁵ One possible explanation is that unmarried participants may have more flexibility and willingness to raise a pet or playing with other people's pets, which may lead them to be at higher risk. However, our data failed to support this hypothesis as we neither observed significant relationship between marital status and pet ownership (pet ownership rates among married and unmarried participants were 12.9% and 13.1% respectively, $p=0.795$), nor collected participants' behaviours with pets. Further studies are therefore warranted.

Given pet ownership is a strong risk factor in this study, its effect may not be fully controlled for in the multivariate analyses, we further estimated the risks of socio-demographic characteristics after stratification by pet ownership. The results revealed that only unmarried pet owners were at higher risk for pet-related injuries after adjustment for other factors, suggesting that unmarried pet owners should be put as the top priority when developing protective countermeasures in future, though the underpinning mechanism is still unclear. Given the statistical power to test significance reduced after the stratification, the relationship between pet-related injuries and other risk factors among pet owners need to be confirmed with a larger sample size in future.

Table 4 Incidence rates of and risk estimations for pet-related injury in the past year (N=84, with p<0.10 in univariate analysis)

	Incidence rate (%)	Univariate analysis		Multivariate Model 1		Multivariate Model 2	
		OR _{unadj}	(95% CI) OR _{adj}	OR _{unadj}	(95% CI) OR _{adj}	OR _{unadj}	(95% CI) OR _{adj}
Gender							
Male (ref.)	1.04	1.00		1.00		1.00	
Female	1.51	1.45	(0.94 to 2.26)	1.73*	(1.02 to 2.95)	1.54	(0.89 to 2.67)
Age (years)							
0–4	1.98	2.79	(0.94 to 8.27)	NA		NA	
5–14	1.31	1.84	(0.83 to 4.08)	NA		NA	
15–24	2.43	3.43*	(1.85 to 6.37)	NA		NA	
25–44	1.35	1.89	(1.05 to 3.40)	NA		NA	
≥45 (ref.)	0.72	1.00		NA		NA	
Marital status							
Married/cohabiting (ref.)	0.85	1.00		1.00		1.00	
Single/divorced/widowed	1.74	2.06*	(1.31 to 3.24)	1.86*	(1.10 to 3.14)	1.80*	(1.04 to 3.10)
Education							
Primary or below (ref.)	0.62	1.00		NA		NA	
Secondary	1.41	2.31	(1.13 to 4.73)	NA		NA	
Postsecondary†	1.63	2.67	(1.25 to 5.69)	NA		NA	
Occupational status							
White-collar job‡ (ref.)	1.70	1.00		NA		NA	
Blue-collar job§	0.42	0.25	(0.06 to 1.02)	NA		NA	
Housewives	0.94	0.55	(0.26 to 1.17)	NA		NA	
Retired	0.32	0.19*	(0.06 to 0.61)	NA		NA	
Students	1.70	1.00	(0.61 to 1.65)	NA		NA	
Others	1.25	0.73	(0.26 to 2.06)	NA		NA	
Monthly household income (HK\$¶)							
<10 000 (ref.)	0.32	1.00		1.00		1.00	
10 000–19 999	1.32	4.20	(1.24 to 14.2)	10.9*	(1.44 to 83.0)	6.85**	(0.89 to 52.7)
20 000–29 999	1.15	3.64	(1.02 to 12.9)	9.21*	(1.17 to 72.4)	6.08**	(0.76 to 48.7)
≥30 000	1.93	6.18*	(1.89 to 20.2)	14.7**	(2.00 to 108.3)	7.60*	(1.01 to 57.1)
Size of living area (ft²‡‡)							
≤400 (ref.)	0.78	1.00		NA		NA	
401–600	0.72	0.92	(0.43 to 1.96)	NA		NA	
601–700	1.74	2.25	(1.08 to 4.69)	NA		NA	
>700	1.88	2.42*	(1.25 to 4.69)	NA		NA	
Living density (ft²/person‡‡)							
<120 (ref.)	0.59	1.00		1.00		1.00	
120–159	1.00	1.70	(0.72 to 3.98)	1.31	(0.51 to 3.37)	1.05	(0.39 to 2.77)
160–249	1.44	2.45	(1.16 to 5.20)	1.89	(0.82 to 4.34)	1.19	(0.50 to 2.86)
≥250	1.65	2.81	(1.27 to 6.23)	2.87*	(1.21 to 6.82)	1.45	(0.58 to 3.61)
Pet ownership							
No (ref.)	0.19	1.00		–		1.00	
Yes	8.56	48.3*	(25.5 to 91.4)	–		52.0***	(22.1 to 122.4)

*Significant difference level at p=0.05 reached in the univariate analysis. **Significant difference level at p=0.01. ***Significant difference level at p=0.001. Bonferroni correction was applied to correct multiple testing. A significance level of $\alpha=0.00417$ (0.05/12) was therefore adopted.

†Post-secondary education included non-degree and degree courses.

‡White-collar job: including manager and administrators, professionals, associate professionals, clerical support workers and service and sales workers.

§Blue-collar job: including craft and related workers, plant and machine operators, elementary occupations and skilled agricultural and fishery workers.

¶Annual exchange rate in 2009: HK\$1=US\$0.129 (US\$1=HK\$7.75); HK\$10 000=US\$1290; HK\$20 000=US\$2580; HK\$30 000=US\$3870.

‡‡1 ft²=0.093 m² (1 m²=10.76 ft²).

NA, the variables were selected but excluded from the Model 1 according to the variable selection method with p<0.10 as the entry and removal criteria; ref., reference group; –, pet ownership was not put in the Model 1.

Most pet-related injuries in our study involved a person's hands, which is consistent with the results from previous studies.^{3 7 18 19} Although injuries examined in this study were not severe, it should be noted that

animals, regardless of the types involved, may carry many diseases that are readily transmissible to human beings, and many of those diseases can be vaccinated against. One such disease is rabies, especially from dog

bites. Rabies is nearly always fatal, and there is no specific treatment.⁸ The microorganism most commonly isolated from wounds by cat bites is *Pasteurella multocida*. Injuries caused by cat bites and scratches may result in cat-scratch disease, which is caused by the bacterium *Bartonella henselae*.^{2 20} Another cat-associated infection is toxoplasmosis, which is caused by the parasite *Toxoplasma gondii* and manifested as a congenital infection.²⁰ It should be noted that *Toxoplasma* infection rates in cat bites have been reported to be between 30–50%, which is much higher than the rate of infection for dog bites (4%).²¹ Unfortunately, only 70.3% of the pets involved in this study had received vaccinations before, with an even lower vaccination rate of 63.6% being found for cats. As cats are one of the most popular types of pets kept in urban households, cat owners need to be educated about the risks of injuries caused by cats and empowered by effective means to protect themselves from such injuries.

Limitations

The results of this study are subject to several limitations. This is a cross-sectional study, thus, it cannot draw a conclusion on the cause–effect relationship. The data on pet-related injuries are self-reported and therefore subject to potential recall bias. This study focuses on pet-related injuries that occurred in a household setting and does not include pet-related injuries that occurred beyond this setting (eg, roads, parks). As the number of participants reporting pet-related injuries was relatively small in this study (n=84), the estimations of the risk factors are inconclusive. It is possible that the associations observed in this study may be just a random phenomenon. Owing to the time constraint, the detailed injury information was collected only from those pet-related injuries which were reported as the most recent injury by the participants. Thus, the results (table 3) may not reflect the real situation of all pet-related injuries in this study. However, as there was no significant difference between the two groups (all pet-related injuries and those as the most recent injury) as shown in table 2, the results may not be biased much from those of all the injuries. Further studies with a prospective longitudinal design are needed to capture all pet-related injuries and confirm the associations observed in this study. Moreover, we did not collect detailed information of pet ownership, for example, dog/cat breeds, history of raising pets, numbers of concurrent pets, the time length of having pets, knowledge and practice on raising pets, all of which are needed to develop effective protection measures and therefore should be addressed in future studies. Finally, some households in this study might keep pets under unlawful circumstances, and therefore participants might be less likely to report owning pets. Thus, the number of pet owners might be under-reported.

CONCLUSIONS

We estimate that more than 86 000 residents in Hong Kong sustained a non-fatal pet-related injury in the household setting in the past 12 months, with a person-based incidence rate of 1.24%. Most of the injuries reported are minor, induced by cats or dogs, with the wounds sustained affecting a person's hands and dealt with by the victims. Pet ownership is a strong risk factor for such an injury. Unmarried and affluent people are also at higher risk, especially unmarried pet owners. Given the increasing trend in pet ownership, it is necessary to take effective measures to protect people from pet-related injuries, especially the unmarried pet owners. Further studies with a prospective longitudinal design should include pet-related injuries that happen outside a household setting.

Author affiliations

¹JC School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong SAR, China

²Department of Physical Education, Hong Kong Baptist University, Hong Kong SAR, China

³Injury Prevention Research Center, Medical College of Shantou University, Shantou, China

Acknowledgements The authors would like to thank Mr Zhe Huang for the research support during the project, Dr Chunlan Guo and Dr Susan Wang for the editing and administrative supports.

Contributors EYYC and YG participated in planning the study, data collection, interpreting the results and final paper writing. LL and PYL participated in data collection, interpreting the results and final paper writing. EYYC obtained research funding. YG wrote the first draft of the article. All authors were involved in planning the article, critical review and editing of the first draft and subsequent revisions to the paper. All authors read and approved the final manuscript.

Funding This project is funded by Health and Health Services Research Fund, Food and Health Bureau, Government of the Hong Kong SAR (grant no. 06070491). The content is solely the responsibility of the authors and does not necessarily represent the official views of the Department of Health, HKSAR.

Competing interests None declared.

Ethics approval Survey and Behavioural Research Ethics Committee at the Chinese University of Hong Kong.

Provenance and peer review Not commissioned; internally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. Lessenger JE. *Agricultural Medicine: a practical guide*. New York, USA: Springer, 2006.
2. Palacio J, León-Artozqui M, Pastor-Villalba E, *et al*. Incidence of and risk factors for cat bites: a first step in prevention and treatment of feline aggression. *J Feline Med Surg* 2007;9:188–95.
3. Centers for Disease Control and Prevention. Nonfatal dog bite-related injuries treated in hospital emergency departments—United States, 2001. *Morb Mortal Wkly Rep* 2003;52:605–10.



4. Stevens JA, Haileyesus T, Centers for Disease Control and Prevention. Nonfatal fall-related injuries associated with dogs and cats--United States, 2001--2006. *Morb Mortal Wkly Rep* 2009;58:277--81.
5. Batson A. *Global companion animal ownership and trade: project summary, June 2008*. 2008. <https://www.zotero.org/groups/chagasepid8515/items/PHEQZ2HV> (accessed 20 Jul 2015).
6. Zhuang J. *China: changing attitudes to pet ownership drive pet food sales*. 2005. <http://blog.euromonitor.com/2005/03/china-changing-attitudes-to-pet-ownership-drive-pet-food-sales.html> (accessed 20 Jul 2015).
7. Ozanne-Smith J, Ashby K, Stathakis VZ. Dog bite and injury prevention--analysis, critical review, and research agenda. *Inj Prev* 2001;7:321--6.
8. Fèvre EM, Kaboyo RW, Persson V, *et al*. The epidemiology of animal bite injuries in Uganda and projections of the burden of rabies. *Trop Med Int Health* 2005;10:790--8.
9. Georges K, Adesiyun A. An investigation into the prevalence of dog bites to primary school children in Trinidad. *BMC Public Health* 2008;8:85.
10. Centre for Health Protection, HKSAR. *Injury Survey 2008*. http://www.chp.gov.hk/files/pdf/injury_survey_eng.pdf (accessed 20 Jul 2015).
11. Census and Statistics Department. *Hong Kong Annual Digest of Statistics: 2012 Edition*. 2012. <http://www.censtatd.gov.hk/> (accessed 13 Mar 2013).
12. Chan EYY, Kim JH, Ng Q, *et al*. A descriptive study of nonfatal, unintentional home-based injury in urban settings: evidence from Hong Kong. *Asia Pacific J Public Health* 2008;20(Suppl):39--48.
13. Lau JTF, Yeung NCY, Choi KC, *et al*. Acceptability of A/H1N1 vaccination during pandemic phase of influenza A/H1N1 in Hong Kong: population based cross sectional survey. *BMJ* 2009;339 (oct27_1):b4164.
14. McGee K, Sethi D, Peden M, *et al*. Guidelines for conducting community surveys on injuries and violence. *Int J Inj Contr Saf Promot* 2004;11:303--6.
15. Chan EYY, Kim JH, Griffiths SM, *et al*. Does living density matter for nonfatal unintentional home injury in Asian urban settings? Evidence from Hong Kong. *J Urban Health* 2009;86:872--86.
16. Woodward M. *Epidemiology: study design and data analysis*. 2nd edn. Boca Raton: Chapman & Hall/CRC, 2005.
17. Thomas N, Brook I. Animal bite-associated infections: microbiology and treatment. *Expert Rev Anti-infe* 2011;9:215--26.
18. Vargo D, DePasquale JM, Vargo AM. Incidence of dog bite injuries in American Samoa and their impact on society. *Hawaii J Med Public Health* 2012;71:6--12.
19. Quirk JT. Non-fatal dog bite injuries in the U.S.A., 2005--2009. *Public Health* 2012;126:300--2.
20. Sandora TJ, Bernstein HH. Neonatal jaundice, animal-induced injuries, and immunizations. *Curr Opin Pediatr* 2001;13:377--85.
21. Smith PF, Meadowcroft AM, May DB. Treating mammalian bite wounds. *J Clin Pharm Ther* 2000;25:85--99.

BMJ Open

Injuries caused by pets in Asian urban households: a cross-sectional telephone survey

Emily Y Y Chan, Yang Gao, Liping Li and Po Yi Lee

BMJ Open 2017 7:

doi: 10.1136/bmjopen-2016-012813

Updated information and services can be found at:
<http://bmjopen.bmj.com/content/7/1/e012813>

These include:

References

This article cites 15 articles, 2 of which you can access for free at:
<http://bmjopen.bmj.com/content/7/1/e012813#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Epidemiology](#) (2145)
[Public health](#) (2262)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>