

3-2010

## Comparison of the anti-inflammatory and anti-nociceptive effects of three medicinal plants known as “Snow Lotus” herb in traditional Uighur and Tibetan medicines

Tao Yi  
*Hong Kong Baptist University, yitao@hkbu.edu.hk*

Zhong-Zhen Zhao  
*Hong Kong Baptist University, zzzhao@hkbu.edu.hk*

Zhi-Ling Yu  
*Hong Kong Baptist University, zlyu@hkbu.edu.hk*

Hu-Biao Chen  
*Hong Kong Baptist University, hbchen@hkbu.edu.hk*

Follow this and additional works at: [https://repository.hkbu.edu.hk/hkbu\\_staff\\_publication](https://repository.hkbu.edu.hk/hkbu_staff_publication)



Part of the [Medicine and Health Sciences Commons](#)

This document is the authors' final version of the published article.

Link to published article: <http://dx.doi.org/10.1016/j.jep.2010.01.037>

---

### APA Citation

Yi, T., Zhao, Z., Yu, Z., & Chen, H. (2010). Comparison of the anti-inflammatory and anti-nociceptive effects of three medicinal plants known as “Snow Lotus” herb in traditional Uighur and Tibetan medicines. *Journal of Ethnopharmacology*, 128 (2), 405-411. <https://doi.org/10.1016/j.jep.2010.01.037>

This Journal Article is brought to you for free and open access by HKBU Institutional Repository. It has been accepted for inclusion in HKBU Staff Publication by an authorized administrator of HKBU Institutional Repository. For more information, please contact [repository@hkbu.edu.hk](mailto:repository@hkbu.edu.hk).

1  
2  
3 **Comparison of the Anti-inflammatory and Anti-nociceptive Effects of Three Medicinal Plants**  
4 **Known as “Snow Lotus” Herb in Traditional Uighur and Tibetan Medicines**  
5  
6

7  
8 Tao Yi, Zhong-Zhen Zhao, Zhi-Ling Yu, Hu-Biao Chen\*  
9

10  
11  
12  
13  
14 **Affiliation**  
15

16 School of Chinese Medicine, Hong Kong Baptist University, Hong Kong Special Administrative Region, P. R. China  
17  
18  
19  
20

21 \* Corresponding author. Tel.: +852 3411 2060; fax: +852 3411 2461.  
22 *E-mail address:* hbchen@hkbu.edu.hk (H.B. Chen).  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 **ABSTRACT**

4 *Ethnopharmacological relevance:*

5 *Saussurea involucreata* (Kar. et Kir.) Sch.-Bip. (Compositae) has long been used under the herbal name  
6 “Snow Lotus” for the treatment of rheumatoid arthritis, stomachache and dysmenorrhea in Uighur folk  
7 medicine. In traditional Tibetan medicine, *S. laniceps* Hand.-Mazz. and *S. medusa* Maxim. have also  
8 been used under the name “Snow Lotus” and prescribed for the treatment of pain and inflammatory  
9 conditions.  
10

11  
12 *Aim of the study:*

13  
14 The present study evaluated the pharmacological effects of three species of “Snow Lotus” in  
15 experimental inflammation and pain models, and determined the chemical compounds that may  
16 correlate with their pharmacological activities.  
17

18  
19 *Material and methods:*

20 The anti-inflammatory activities of the three herbs were observed by using carrageenan-induced paw  
21 edema in rats and xylene-induced ear edema in mice. Investigations on the analgesic effects were  
22 conducted, including acetic acid-induced writhing and hot-plate test. An UPLC-MS method was  
23 developed to analyse the chemical composition of the three herbs and of plasma samples after herb  
24 administration.  
25

26  
27 *Results:*

28 In rat paw edema model, the peak inhibitory effects of *S. laniceps* and *S. involucreata* (55.1% and 42.2%,  
29 respectively) were recorded with the dose of 400 mg/kg at 3 h post carrageenan injection. In mouse ear  
30 edema model, oral administration of *S. laniceps*, *S. involucreata* and *S. medusa* extract (400 mg/kg)  
31 resulted in a significant inhibition of ear edema by 40.9%, 33.3%, and 9.1%, respectively. In the  
32 writhing test, oral administration of *S. laniceps* extract (100, 200 and 400 mg/kg) resulted in a  
33 significant inhibition of writhings by 13.5%, 22.3%, and 43.5%, respectively. In the hot plate test, *S.*  
34 *laniceps* extract significantly increased the latency of jumping response by 38.2% and 52.7% when  
35 treated orally at 200 and 400 mg/kg in mice, respectively. Flavonoids, coumarins and lignins were  
36 found to be present in plasma after administration of the extracts and may be on the basis of the  
37 observed pharmacological effects.  
38  
39  
40

41  
42 *Conclusion:*

43 The results clearly demonstrated that *S. laniceps* was most effective; *S. involucreata* exhibited a  
44 moderate potency, whereas *S. medusa* possessed little effect against the experimental edema and pains.  
45 This study also supported discrimination among the three herbs when using them in folk medicine.  
46

47  
48 *Keywords:*

49 Snow Lotus; *Saussurea involucreata*; *Saussurea laniceps*; *Saussurea medusa*; Anti-inflammatory;  
50 Anti-nociceptive  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 **1. Introduction**  
4

5 Because of biodiversity, related medicinal plants from the same family or genus have been and are  
6 being used for similar therapeutic purposes in various folk medicines. *Saussurea involucrata* (Kar. et  
7 Kir.) Sch.-Bip. (SI, Fig. 1), of the Composite family, grows in the mountains at heights of 4000-4300 m  
8 in the Tianshan and A'er Tai areas in China (Chen et al., 1999). With a reputation for diminishing  
9 inflammation and facilitating blood circulation, the dried aerial parts of SI have long been used under  
10 the name "Snow Lotus" for the treatment of rheumatoid arthritis, stomachache and dysmenorrhea in  
11 Uighur folk medicine (Chinese Pharmacopoeia Commission, 2005). Two other species of the same  
12 genus, *S. laniceps* Hand.-Mazz. and *S. medusa* Maxim. (SL and SM, Fig. 1), mainly distributed in the  
13 Qinghai-Tibet plateau at heights of 3500-5300 m, have also been used under the name "Snow Lotus"  
14 and prescribed for the treatment of pain and inflammatory conditions in Tibetan folk medicine  
15 (Commission of Chinese Ethnomedicine, 1984).  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

26 *Insert Fig. 1 here*  
27  
28

29 Currently, these three species of *Saussurea* genus are all used as an herbal drug for the treatment of  
30 inflammation and pain-related diseases in Chinese medicine (Commission of Chinese Materia Medica,  
31 1999). However, whether these related plants can be used interchangeably is still unknown thereby  
32 limiting further bioactive screening and clinical popularization. This is particularly important because  
33 one of the species, SI, is in danger of extinction due to the pressures of herbal collection (Ministry of  
34 Environmental Protection of the People's Republic of China, 1987). Studies on the pharmacological  
35 effects of the three medicinal plants have been performed using various models (Zheng et al., 1993;  
36 Duan et al., 2002; Gao et al., 2005); however, the existing studies have each focused on the  
37 pharmacological investigation of a single herb, and no systematic comparison of the pharmacological  
38 potencies related to the clinical effects of these herbs has been reported. Therefore, to encourage  
39 reasonable and sustainable use of these related species, a comparison of their chemical composition and  
40 pharmacological activities is urgently needed.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

51 In our previous study, we reported the chemical composition of these three herbs used under the  
52 name "Snow Lotus" (Yi et al., 2009a; Yi et al., 2009b). In this follow-up study, we compare the  
53 suppressive effects of the three herbs in two inflammatory models and two nociceptive models in  
54 rodents. In addition we sought to determine the correlation between herb constituents absorbed in the  
55 blood stream and pharmacological activity. To do this, the herbal constituents found in blood plasma  
56 after herb administration were detected using ultra-performance liquid chromatography-electrospray  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 ionization mass spectrometry (UPLC-ESI-MS). The results reveal that the pharmacological potencies  
4 of the three herbs varied, which would contribute to the different chemical composition and absorbed  
5 constituents of the three herbs. As for physiological effects, among the three herbs, SL significantly  
6 inhibited inflammatory responses and nociceptive production to a greater degree than either of the other  
7 two. Thus, SL is the most likely candidate to become a complementary and alternative drug for the  
8 treatment of inflammatory and painful diseases.  
9

## 16 **2. Materials and methods**

### 17 *2.1 Plant materials and reagents*

20 *Saussurea involucrata* (Kar. et Kir.) Sch.-Bip. (SI) was collected from the Tianshan areas in the  
21 Xinjiang Uighur Autonomous Region of China in 2007. *S. laniceps* Hand.-Mazz. (SL) and *S. medusa*  
22 Maxim. (SM) were collected from Lhasa in Tibet of China in 2007 and 2005, respectively. Identity of  
23 the herbs was confirmed by Dr. Hubiao Chen (School of Chinese Medicine, Hong Kong Baptist  
24 University, Hong Kong) by means of geographical origin and macroscopic character assessment.  
25 Voucher specimens were deposited in the Chinese Medicines Center, Hong Kong Baptist University  
26 (TS-03 for SI, MT-01 for SL and SM-04 for SM, respectively).  
27  
28  
29  
30  
31  
32

33 Indomethacin, carrageenan, acetic acid and carboxy methylcellulose were purchased from Sigma  
34 Chemical Co. (St. Louis, MO, USA). Rotundine, an analgesic drug was purchased from Zhenang  
35 Pharmaceutical Co., Ltd. (Nanjing, China). Acetonitrile and methanol of chromatography grade were  
36 purchased from Lab-scan (Bangkok, Thailand). Formic acid of chromatography grade and ethanol of  
37 analytical grade were purchased from Merck (Darmstadt, Germany). Water was purified using a  
38 Milli-Q water system (Millipore; Bedford, MA, USA).  
39  
40  
41  
42  
43  
44

### 45 *2.2 Animals*

47 ICR mice weighing 18-23 g and SD rats weighing 150-200 g were purchased from the Laboratory  
48 Animal Services Center, the Chinese University of Hong Kong, Hong Kong. All animals were fed with  
49 a standard chow diet and water. All animals were housed at a room temperature of  $23 \pm 1$  °C with a 12  
50 h light/dark cycle. A standard rodent diet and water were provided *ad libitum*. All experimental  
51 protocols were approved by the Committee on the Use of Human & Animal Subjects in Teaching and  
52 Research of Hong Kong Baptist University, in accordance with the Animals Ordinance (Department of  
53 Health, Hong Kong).  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

### 2.3 Preparation of herbal extracts

Plant materials were cut into small pieces and ground into powder. Herbal sample powder (0.5 kg) was extracted with 50% ethanol by means of percolation at room temperature. The operations were repeated until the extract became colorless. The combined extracts were evaporated to remove ethanol at reduced pressure in a rotary evaporator (50 °C), and then were lyophilized with a freeze-dry system. SI extract (129 g, yield 25.8%, w/w), SL extract (86 g, yield 17.2%, w/w) and SM extract (56 g, yield 11.2%, w/w) were thus obtained. The dried extracts were suspended in 1% (w/v) aqueous carboxy methylcellulose for administration to animals.

### 2.4 Carrageenan-induced paw edema in rat

For the determination of effects on acute inflammation, the carrageenan-induced paw edema model described by Yesilada and Kupeli (Yesilada and Kupeli, 2002) was employed with modifications. SD rats were weighed and randomized into eleven groups of ten animals each, namely a control group (aqueous carboxy methylcellulose-treated), a reference drug group (indomethacin-treated), and three groups each (receiving 100, 200, or 400 mg/kg of drug) for SI treatment, SL treatment and SM treatment. The test agents were orally administered to the rats for 5 consecutive days. One hour after the last delivery, 100 µl of a 1% solution of carrageenan in 0.9% physiological saline was injected subcutaneously into the subplantar region of the right hind paw. The left hind paw received 100 µl of saline solution similarly injected. The difference in footpad volume between the right and left foot was measured with a water displacement plethysmometer (plethysmometer 7150, UGO Basile, Italy) at 1, 3, and 5 h after induction of inflammation. Mean differences of treated groups were compared with the mean difference of the control group and analysed using statistical methods.

### 2.5 Xylene-induced ear edema in mice

The xylene-induced ear edema test with modifications was performed as previously described (Luo et al., 2008). Oral administrations with vehicle, reference drug and herbal extracts before inducing ear edema were conducted for 5 consecutive days. Thirty minutes after the last administration of tested drugs, a total of 20 µl of xylene was applied to the inner and outer surface of the right ear of each mouse. The left ear remained untreated. Control animals received the irritant and an equal volume of aqueous carboxy methylcellulose, while indomethacin (10 mg/kg) served as the reference. Fifteen minutes later, the mice were sacrificed by cervical dislocation and the plugs were removed with a cork borer (6.5 mm in diameter) from both the right and left ear. The difference in weight between the two plugs was taken as a measure of edematous response.

## 2.6 Writhing test in mice

Writhing test was performed as the chemical pain model (Arihan et al., 2009). Briefly, vehicle, rotundine (50 mg/kg), and test solution (100, 200 and 400 mg/kg) were orally administered for five consecutive days. Half hour after the last delivery, 0.2 ml of acetic acid (0.7% w/v) in saline was injected intraperitoneally. The numbers of abdominal writhing movements were recorded for 15 min starting 5 min after *i.p.* injection in each animal.

## 2.7 Hot plate test in mice

The hot plate test was used as the thermal pain model (Suzuki et al., 2009). Briefly, vehicle, rotundine (50 mg/kg), and test solution (100, 200 and 400 mg/kg) were orally administered for five consecutive days. Half hour after the last delivery, each mouse was placed on a hot-plate surface (IITC Model 39, Woodland Park, CA) maintained at  $55 \pm 0.2^\circ\text{C}$ . The reaction time from hot-plate placement to hind-paw lick was recorded. The cut-off time was set at 60 second to prevent tissue injury. Before the experiments, all animals were tested for heat stimulation latency, and those with response time  $< 5$  second or  $> 30$  second were excluded.

## 2.8 UPLC-MS fingerprint analysis of herb extracts and animal plasma

A Waters Acquity<sup>TM</sup> ultra-performance liquid chromatography (UPLC) system (Waters Corp., Milford, USA) consisting of a binary pump, autosampler, thermostated column compartment and diode array detector (DAD), was used for fingerprint analysis. For mass spectrometric determination, the UPLC system was hyphenated to a Bruker MicrOTOFQ system by an electrospray ionization (ESI) interface (Bruker Daltonics, Bremen, Germany).

The chromatographic separation was carried out on a Waters BEH C<sub>18</sub> column (1.7  $\mu\text{m}$ , 2.1  $\times$  100 mm, Waters Corp.) with a VanGuard<sup>TM</sup> pre-column (BEH, C<sub>18</sub>, 1.7  $\mu\text{m}$ , 2.1  $\times$  5 mm). The mobile phase consisted of 0.1% formic acid in water (A) and 0.1% formic acid in acetonitrile (B) using a gradient program of 3% (B) in 0-2 min, 3-15% in 2-14 min, 15-20% (B) in 14-22 min and 20-45% (B) in 22-30 min. The solvent flow rate was 0.3 ml/min, the column temperature was set to 40  $^\circ\text{C}$  and the detection wavelength was 280 nm. The conditions of MS analysis in the positive ion mode were as follows: drying gas (nitrogen), flow rate, 8 l/min; gas temperature, 200  $^\circ\text{C}$ ; scan range, 50-1000 *m/z*; end plate offset voltage, -500 V; capillary voltage, 4500 V; nebulizer press, 1.5 Bar.

For analysis of herb extracts, three extracts of 0.1 g each were dissolved in 50 ml of methanol, and then centrifuged at 12000 rpm for 10 min to obtain sample solution. For the analysis of animal plasma

1  
2  
3 after herb administration, the three herbal extracts (400 mg/kg) were orally administered to three  
4 groups of two rats each for 5 consecutive days. Three hours after the last delivery, blood samples (ca.  
5 0.6 ml) were withdrawn from the fossa orbitalis into heparin-stabilized test-tubes and immediately  
6 centrifuged at 3000 rpm for 5 min to obtained plasma. One milliliter methanol was added into 0.2 ml  
7 plasma and vortex-mixed for 5 s, then centrifuged at 12000 rpm for 10 min to precipitate the protein  
8 and extracted analytes. The supernatant was transferred into another tube and dried under a stream of  
9 nitrogen. The residue was reconstituted with 0.1ml methanol and then centrifuged at 12000 rpm for 10  
10 min. To obtain mouse plasma after drug administration, the three herbal extracts (400 mg/kg) were  
11 orally administered to three groups of five mice each for 5 consecutive days, respectively. One hour  
12 after the last delivery, the mice were sacrificed by decapitation, and the blood was collected and  
13 handled using the procedures described above. An aliquot of 3  $\mu$ l for the supernatant of all the test  
14 solutions was analyzed by UPLC–MS.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

## 26 *2.9 Statistical analysis*

27  
28 Values obtained from experiments were expressed as mean  $\pm$  S.E.M and further analyzed using  
29 one-way ANOVA followed by Dunnett test for multiple comparisons, with the level of significance  
30 chosen as  $P < 0.05$ .  
31  
32  
33

## 34 **3. Results and discussion**

### 35 *3.1 Suppressive effects in inflammatory models*

36  
37  
38 Carrageenan-induced rat paw edema and xylene-induced mouse ear edema are the typical animal  
39 models used to evaluate the anti-inflammatory effect of natural products; they have a good reputation  
40 for screening anti-inflammatory agents. Inflammation induced by carrageenan is an acute and highly  
41 reproducible inflammatory model. Cardinal signs of inflammation develop immediately following  
42 subcutaneous injection, resulting from action of proinflammatory agents. The inflammatory response is  
43 usually quantified by increase in paw size (edema) and is also modulated by inhibitors within the  
44 inflammatory cascade (Winyard and Willoughby, 2003).  
45  
46  
47  
48  
49  
50  
51

52 In our study, the suppressive effects of the three herbal extracts on carrageenan-induced paw edema  
53 in rat are shown in Table 1. The extracts of the three herbs exhibited varying degrees of  
54 anti-inflammatory activity. SL extract administered orally at doses of 100-400mg/kg showed significant  
55 dose-dependent reduction of the carrageenan-induced paw edema, SI extract showed a moderate  
56 inhibition of edema formation, whereas SM extract apparently had little effect on edema. The peak  
57 inhibitory effects of SL and SI (55.1% and 42.2%, respectively) were recorded with the dose of 400  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3 mg/kg at 3 h post-carrageenan injection ( $p < 0.01$ ), compared with the indomethacin-treated group (10  
4 mg/kg, 75.5%).

5  
6  
7 *Insert Table 1 here*

8  
9 Xylene-induced mouse ear edema leads to an acute inflammatory response characterized by fluid  
10 accumulation and edema in the *in vivo* model. Suppression of this response is taken as an indication of  
11 antiphlogistic effect (Atta and Alkofahi, 1998). The suppressive effects of the three herbal extracts on  
12 xylene-induced mouse ear edema are shown in Figure 2. In our study, SL extract administered orally at  
13 a dose of 200-400mg/kg and SI extract at a dose of 400mg/kg significantly reduced ear edema, and the  
14 mean weight of ear edema in SL-treated group (3.9 mg) of 400 mg/kg was lower than that of the  
15 SI-treated group (4.4 mg) as well as the SM-treated group (6.0 mg) at equivalent doses. Compared with  
16 the control group (6.6 mg), oral administration of SL, SI and SM extract (400 mg/kg) resulted in a  
17 significant inhibition of ear edema by 40.9%, 33.3%, and 9.1%, respectively.

18 Hence, SL extract appeared to be the most effective of the three herbs in reducing inflammation,  
19 implying that SL possesses anti-inflammatory components.

20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31 *Insert Fig. 2. here*

### 32 33 3.2 *Suppressive effects in analgesic models*

34  
35 Animal pain tests have been developed primarily for the screening of potential analgesic drugs.  
36 Various tests have different profiles of sensitivity; thus, while no single laboratory test procedure can  
37 give a comprehensive evaluation of analgesics, a combination of several tests provides a good  
38 substitute (Boulton et al., 1989). Therefore, the methods of both peripherally and centrally mediated  
39 effects were selected for investigating anti-nociceptive effect. The acetic acid-induced abdominal  
40 constriction and hot-plate methods are the animal models typically used to elucidate peripheral and  
41 central activity, respectively (Verma et al., 2005).

42  
43 In the acetic acid-induced abdominal writhing test (Table 2), oral administration of SI (100, 200 and  
44 400 mg/kg) resulted in a significant inhibition of writhing by 12.4%, 29.5%, and 35.2%, respectively,  
45 and oral administration of SL (100, 200 and 400 mg/kg) resulted in a significant inhibition of writhings  
46 by 13.5%, 22.3%, and 43.5%, respectively. The SI and SL extract at a dose of 400 mg/kg inhibited the  
47 acetic acid-induced pain with a potency comparable to that of the reference drug, rotundine ( $p > 0.05$ ).  
48 Acetic acid itself may cause pain; at the same time, it can also stimulate the tissue to produce several  
49 mediators such as histamine, serotonin, cytokines, and eicosanoids with an increase in peritoneal fluid  
50 levels of these mediators (Bentley et al., 1983, Rinaldi et al., 2009). Therefore, anti-nociceptive activity

1  
2  
3 of SI and SL may be related to the reduction in the liberation of those inflammatory mediators or by  
4 direct blockage of receptors resulting in peripheral anti-nociceptive effects.

5  
6 To check for possible central anti-nociceptive activity of the herbs, the hot plate test was performed,  
7 and the results are shown in Table 2. When mice were treated with extracts of SL (100, 200 and 400  
8 mg/kg), there was a dose-dependent increase in response to thermal stimulation compared with control  
9 mice. These results clearly indicate that SL significantly increased the latency of jumping response by  
10 38.2% and 52.7% when treated at 200 and 400 mg/kg, respectively, suggesting that SL has central  
11 analgesic properties.  
12  
13  
14  
15  
16

17 *Insert Table 2 here*

### 18 19 20 *3.3 Phytochemicals and absorbed constituents after herb administration*

21  
22 Ultra-performance liquid chromatography (UPLC), a new category of analytical separation science,  
23 retains the practicality and principles of high-performance liquid chromatography (HPLC) while  
24 increasing the overall interlaced attributes of speed, sensitivity, and resolution. In the present study, the  
25 extracts of the three herbs and plasma samples after herb administration were analyzed using  
26 UPLC-MS, and the typical chromatograms are shown in Figure 3 to Figure 5. Based on our previous  
27 study (Yi et al., 2009a; Yi et al., 2009b), the main constituents of the three herbs are found to be:  
28 organic acids and flavonoids in SI, coumarins and their glucosides in SL, and lignins in SM.  
29  
30  
31  
32  
33  
34

35 For the analysis of absorbed constituents in the SI-treated group (Figure 3), chlorogenic acid, one of  
36 the main constituents in SI extract, was not detected, neither in mouse nor rat plasma samples. In  
37 accordance with the literature, chlorogenic acid possesses antioxidant as well as antiviral effects with  
38 lower bioavailability by oral administration (Olthof et al., 2001; Gonthier et al., 2003). In contrast, rutin,  
39 another main constituent of the SI extract, was detected in mouse plasma. It has been reported that rutin  
40 posses anti-inflammatory and anti-nociceptive effects (Jia et al., 2005). Therefore, the moderate  
41 inhibition of SI on the inflammatory and nociceptive models in this study would contribute to the  
42 presence of rutin in SI herb.  
43  
44  
45  
46  
47  
48  
49

50 *Insert Fig. 3. here*

51  
52 For the analysis of absorbed constituents in the SL-treated group, umbelliferone and scopoletin  
53 were detected both in mouse and rat plasma samples. The typical UPLC chromatogram is shown in  
54 Figure 4. The results indicate that umbelliferone and scopoletin are absorbed by animals after SL  
55 extract oral administration. In recent years, it has been reported that umbelliferone and scopoletin are  
56 easy to transport through the small intestine mucosal barrier directly, and their bioavailability compared  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 to inderal, an easily absorbed reference drug (Yang et al., 2008). On the other hand, umbelliferone and  
4 scopoletin showed cyclooxygenase-2 (COX-2) and 5-lipoxygenase (5-LOX) dual inhibitory activity  
5 (Kim et al., 2006), which suggests that the two compounds produce anti-inflammatory activity in part  
6 via the inhibition of the generation of eicosanoids. It has also been reported that umbelliferone and  
7 scopoletin exhibit significant and dose-related antinociceptive effects against acetic acid-induced  
8 visceral pain in mice (Tanaka et al., 1977; Meotti et al., 2006). Moreover, scopoletin is a commonly  
9 used anti-asthmatic in modern medicine, and it is mainly prescribed for the treatment of acute and  
10 chronic bronchitis. Therefore, the anti-inflammatory and analgesic effect of SL extract in the present  
11 study would contribute to the relatively greater bioavailability as well as the pharmacological activities  
12 of umbelliferone and scopoletin.  
13  
14  
15  
16  
17  
18  
19  
20

21 *Insert Fig. 4. here*  
22

23 For the analysis of the absorbed constituents in the SM-treated group (Fig. 5), arctigenin, an  
24 aglycone and the main metabolite of arctiin (Zheng et al., 2005), was detected in rat plasma samples.  
25 However, the pharmacological activities of arctigenin are mainly anti-cancer and anti-viral (Awale et  
26 al., 2006; Wang et al., 2006); they are not reported to have strong anti-inflammatory activity. Thus,  
27 these features are probably related to the small suppressive effects of SM on inflammation and  
28 nociception in the models in the present study. It is speculated that SM and SL have been and are being  
29 interchangeably used in Tibetan folk medicine because of their taxonomic similarity rather than  
30 equivalence in pharmacological effectiveness.  
31  
32  
33  
34  
35  
36  
37  
38

39 *Insert Fig. 5. here*  
40

#### 41 **4. Conclusion**

42 The present study investigated the pharmacological effects of three species traditionally used under  
43 the herbal drug name “Snow Lotus” against experimental inflammation and pain. Four animal models,  
44 i.e., carrageenan-induced rat paw edema and xylene-induced mouse ear edema models, were used to  
45 study anti-inflammatory activity, and acetic acid-induced mice abdominal writhing and mouse hot-plate  
46 models were used to study the anti-nociceptive effect. All animal models clearly demonstrated that  
47 *Saussurea laniceps* was most effective; *S. involucrata* exhibited a moderate potency, whereas *S.*  
48 *medusa* possessed little effect against the experimental edema and pains.  
49  
50  
51  
52  
53  
54  
55

56 Comparative analysis of the three herbs and of blood plasma after herb administration was used to  
57 elucidate correlation between compounds and pharmacological activity. The results revealed that the  
58 chemical composition, absorbed constituents and pharmacological activities of the three herbs were  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

different, which supported discrimination among the three herbs when using them in folk medicine. The results demonstrate that *S. laniceps* is the most potent of the three, and suggested that it is the best species to be used in traditional medicine as an anti-inflammatory and analgesic drug instead *S. involucrata* or *S. medusa*. The results also indicate that *S. involucrata* is less potent than *S. laniceps*; if this information was disseminated among herb collectors and drug dealers it could help reduce collection pressures on this rare and endangered species.

### **Acknowledgements**

This research was funded by the Faculty Research Grant of Hong Kong Baptist University (FRG/07-08/II-38).

## References

- Arihan, O., Boz, M., Iskit, A.B., Ilhan, M., 2009. Antinociceptive activity of coniine in mice. *Journal of Ethnopharmacology* 125, 274–278.
- Atta, A.H., Alkofahi, A., 1998. Anti-nociceptive and anti-inflammatory effects of some Jordanian medicinal plant extracts. *Journal of Ethnopharmacology* 60, 117–124.
- Awale, S., Lu, J., Kalauni, S.K., Kurashima, Y., Tezuka, Y., Kadota, S., Esumi, H., 2006. Identification of arctigenin as an antitumor agent having the ability to eliminate the tolerance of cancer cells to nutrient starvation. *Cancer Research* 66, 1751–1757.
- Bentley, G.A., Newton, S.H., Starr, J., 1983. Studies on the anti-nociceptive action of against drugs and their interaction with opioid mechanisms. *British Journal of Pharmacology* 79, 125–134.
- Boulton, A.A., Baker, G.B., Greenshaw, A.J., 1989. *Psychopharmacology*. Humana Press, New Jersey, 145–216.
- Chen, F.J., Yang, Y.G., Zhao, D.X., Gui, Y.L., Guo, Z.C., 1999. Advances in studies of species, habitats distribution and chemical composition of snow lotuses (*Saussurea*) in China. *Zhiwuxue Tongbao (Chinese Bulletin of Botany)* 16, 561–566.
- Chinese Pharmacopoeia Commission, 2005. Chinese Pharmacopoeia Commission, Pharmacopoeia of the People's Republic of China (2005) vol. 1. Chemical Industry Publishing Press, Beijing, China, pp. 36–37.
- Commission of Chinese Ethnomedicine, 1984. *Ethnomedicine of China* vol. 1. The Ethnic Publishing House, Beijing, China, pp. 261–263.
- Commission of Chinese Materia Medica, 1999. *Materia Medica of China* vol. 7. Shanghai Scientific and Technical Publishers, Shanghai, China, pp. 931–934.
- Duan, H.Q., Takaishi, Y., Momota, H., Ohmoto, Y., Taki, T., 2002. Immunosuppressive constituents from *Saussurea medusa*. *Phytochemistry* 59, 85–90.
- Gao, X., Zhang, Z.M., Xu, A.X., Lei, X.Y., 2005. Antioxidation of ethanolic extract from *Saussurea laniceps* on tissues and erythrocytes in rats. *Zhongguo Yao Xue Za Zhi (Chinese Pharmaceutical Journal)* 40, 1062–1065.
- Gonthier, M.P., Verny, M.A., Besson, C., Remesy, C., Scalbert, A., 2003. Chlorogenic acid bioavailability largely depends on its metabolism by the gut microflora in rats. *Journal of Nutrition* 133, 1853–1859.
- Jia, J.M., Wu, C.F., Liu, W., Yu, H., Hao, Y., Zheng, J.H., Ji, Y.R., 2005. Anti-inflammatory and analgesic activities of the tissue culture of *Saussurea involucrata*. *Biological and Pharmaceutical Bulletin* 28, 1612–1614.
- Kim, J.S., Kim, J.C., Shim, S.H., Lee, E.J., Jin, W.Y., Bae, K., Son, K.H., Kim, H.P., Kang, S.S., Chang, H.W., 2006. Chemical constituents of the root of *Dystaenia takeshimana* and their anti-inflammatory activity. *Archives of Pharmacal Research* 29, 617–623.
- Luo, P., Zhang, Z.F., Yi, T., Zhang, H., Liu, X.F., Mo, Z.J., 2008. Anti-inflammatory activity of the extracts and fractions from *Erigeron multiradiatus* through bioassay-guided procedures. *Journal of Ethnopharmacology* 119, 232–237.
- Meotti, F.C., Ardenghi, J.V., Pretto, J.B., Souza, M.M., Moura, J.A., Junior, A.C., Soldi, C., Pizzolatti, M.G., Santos, A.R.S., 2006. Antinociceptive properties of coumarins, steroid and dihydrostyryl-2-pyrones from *Polygala sabulosa* (Polygalaceae) in mice. *Journal of Pharmacy and Pharmacology* 58, 107–112.
- Ministry of Environmental Protection of the People's Republic of China, 1987. *List of rare and endangered plants for protection in China* vol. 1. Science Press, Beijing, China, pp. 1–96.
- Olthof, M.R., Hollman, P.C.H., Katan, M.B., 2001. Chlorogenic acid and caffeic acid are absorbed in humans. *Journal of Nutrition* 131, 66–71.
- Rinaldi, S., Silva, D.O., Bello, F., Alviano, C.S., Alviano, D.S., Matheus, M.E., Fernandes, P.D., 2009. Characterization of the antinociceptive and anti-inflammatory activities from *Cocos nucifera* L. (Palmae). *Journal of Ethnopharmacology* 122, 541–546.
- Suzuki, Y., Yuzurihara, M., Hibino, T., Yano, S., Kase, Y., 2009. Aqueous extract of *Asiasari radix* inhibits formalin-induced hyperalgesia via NMDA receptors. *Journal of Ethnopharmacology* 123, 128–133.
- Tanaka, S., Ikeshiro, Y., Tabata, M., Konoshima, M., 1977. Anti-nociceptive substances from the roots of *Angelica acutiloba*. *Arzneimittel-Forschung* 27, 2039–2045.
- Verma, P.R., Joharapurkar, A.A., Chatpalliwar, V.A., Asnani, A.J., 2005. Antinociceptive activity of alcoholic extract of *Hemidesmus indicus* R.Br. in mice. *Journal of Ethnopharmacology* 102, 298–301.
- Wang, X.Y., Jia, W., Zhao, A.H., Wang, X.R., 2006. Anti-influenza agents from plants and traditional Chinese medicine. *Phytotherapy Research* 20, 335–341.
- Winyard, P.G., Willoughby, D.A., 2003. *Inflammation protocols*. Humana Press, New Jersey, 126–185.
- Yang, X.W., Guo, Q.M., Wang, Y., 2008. Absorption and transport of 6 coumarins isolated from the roots of *Angelica*

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

*pubescens* f. *biserrata* in human Caco-2 cell monolayer model. *Zhong Xi Yi Jie He Xue Bao (Journal of Chinese Integrative Medicine)* 6, 392–398.

Yesilada, E., Kupeli, E., 2002. *Berberis crataegina* DC. root exhibits potent anti-inflammatory, analgesic and febrifuge effects in mice and rats. *Journal of Ethnopharmacology* 79, 237–248.

Yi, T., Chen, H.B., Zhao, Z.Z., Jiang, Z.H., Cai, S.Q., Wang, T.M., 2009. Identification and determination of the major constituents in the traditional Uighur medicinal plant *Saussurea involucrata* by LC-DAD-MS. *Chromatographia* 69, 537–542.

Yi, T., Chen, H.B., Zhao, Z.Z., Jiang, Z.H., Cai, S.Q., Wang, T.M., 2009. Comparative analysis of the major constituents in the traditional Tibetan medicinal plants *Saussurea laniceps* and *S. medusa* by LC-DAD-MS. *Chromatographia* 70, 957–962.

Zheng, R.L., Liu, G.S., Xing, G.X., Jia, Z.J., Du, M., Tan, L.Q., 1993. Free radical scavenging and anti-fatigue activities of *Saussurea involucrata* polysaccharides. *Acta Pharmacologica Sinica* 14, S47–49.

Zheng, Y.M., Cai, S.X., Xu, X.Y., Fu, S.Q., 2005. Dynamical studies on metabolic chemistry of lignanoids from seeds of *Arctium lappa*. *Zhongguo Zhong Yao Za Zhi (China Journal of Chinese Materia Medica)* 30, 1287–1289.

## Legends for Tables

### **Table 1**

Anti-inflammatory effect of three herbal extracts on carrageenan-induced paw edema in rats.

### **Table 2**

Antinociceptive effect of three herbal extracts on acetic acid-induced writhing and hot plate test in mice.

**Table 1**

Anti-inflammatory effect of three herbal extracts on carrageenan-induced paw edema in rats

Groups	Dose (mg/kg)	Edema volume (EV, ml) and edema inhibition (EI, %) after injection (h)					
		1 h		3 h		5 h	
		EV (ml)	EI (%)	EV (ml)	EI (%)	EV (ml)	EI (%)
Control	-	0.645 ± 0.042	-	0.787 ± 0.048	-	0.756 ± 0.065	-
SI	100	0.528 ± 0.040	18.1	0.636 ± 0.042	19.2	0.625 ± 0.026	17.3
	200	0.572 ± 0.043	11.3	0.604 ± 0.052	23.2	0.555 ± 0.071*	26.6
	400	0.518 ± 0.072	19.7	0.455 ± 0.047**	42.2	0.491 ± 0.035**	35.1
SL	100	0.463 ± 0.032*	28.2	0.541 ± 0.065*	31.3	0.628 ± 0.057	16.9
	200	0.509 ± 0.040	21.1	0.405 ± 0.047**	48.5	0.440 ± 0.041**	41.8
	400	0.469 ± 0.026*	27.3	0.353 ± 0.057**	55.1	0.372 ± 0.040**	50.8
SM	100	0.627 ± 0.044	2.8	0.726 ± 0.065	7.8	0.688 ± 0.048	9.0
	200	0.611 ± 0.043	5.3	0.672 ± 0.053	14.6	0.698 ± 0.025	7.7
	400	0.505 ± 0.060	21.7	0.543 ± 0.061*	31.0	0.630 ± 0.063	16.7
Indomethacin	10	0.226 ± 0.026**	65.0	0.193 ± 0.029**	75.5	0.214 ± 0.040**	71.7

SI: *Saussurea involucrata*; SL: *Saussurea laniceps*; SM: *Saussurea medusa*.Values of edema shown are mean ± S.E.M. ( $n=10$ ).\*  $p < 0.05$ ; \*\*  $p < 0.01$  vs. control group.



**Table 2****Antinociceptive effect of three herbal extracts on acetic acid-induced writhing and hot plate test in mice**

Groups	Dose (mg/kg)	Frequency (Counts/15 min)	Inhibition (%)	Latency time (sec)	Increase (%)
Control	-	19.3 ± 1.2	-	16.5 ± 1.2	-
SI	100	16.9 ± 2.6	12.4	17.4 ± 1.5	5.5
	200	13.6 ± 1.4	29.5	19.2 ± 1.8	16.4
	400	12.5 ± 1.2*	35.2	18.8 ± 1.3	13.9
SL	100	16.7 ± 1.9	13.5	18.3 ± 2.1	10.9
	200	15.0 ± 2.4	22.3	22.8 ± 1.2*	38.2
	400	10.9 ± 1.3**	43.5	25.2 ± 1.5**	52.7
SM	100	18.2 ± 1.3	5.7	17.1 ± 0.9	3.6
	200	19.9 ± 2.1	-3.1	17.9 ± 1.9	8.5
	400	15.4 ± 1.8	20.2	20.1 ± 1.3	21.8
Rotundine	50	8.1 ± 0.8**	58.0	27.7 ± 1.6	67.9

SI: *Saussurea involucrata*; SL: *Saussurea laniceps*; SM: *Saussurea medusa*.

Values of edema shown are mean ± S.E.M. (n=10).

\*  $p < 0.05$ ; \*\*  $p < 0.01$  vs. control group.

## Legends for Figures


**Fig. 1.** Photos of *Saussurea involucrata* (Kar. et Kir.) Sch.-Bip. (SI), *S. laniceps* Hand.-Mazz. (SL), *S. medusa* Maxim. (SM) plants and their medicinal materials.

**Fig. 2.** Anti-inflammatory effect of three herbal extracts on xylene-induced ear edema in mice.


**Fig. 3.** UPLC chromatogram of a sample of SI extract (SI-E), a plasma sample of SI-treated rat (SI-R) and a plasma sample of SI-treated mouse (SI-M). Peak 1 to peak 3 were identified as chlorogenic acid, rutin and 1,5-dicaffeoylquinic acid, respectively.

**Fig. 4.** UPLC chromatogram of a sample of SL extract (SL-E), a plasma sample of SL-treated rat (SL-R) and a plasma sample of SL-treated mouse (SL-M). Peak 1 to peak 7 were identified as umbelliferone 7-*O*- $\beta$ -D-glucoside, chlorogenic acid, scopoletin 7-*O*- $\beta$ -D-glucoside, syringoside, umbelliferone, scopoletin and 1,5-dicaffeoylquinic acid, respectively.


**Fig. 5.** UPLC chromatogram of a sample of SM extract (SM-E), a plasma sample of SM-treated rat (SM-R) and a plasma sample of SM-treated mouse (SM-M). Peak 1 to peak 3 were identified as chlorogenic acid, arctiin and arctigenin, respectively.




**Fig. 1.** Photos of *Saussurea involucrata* (Kar. et Kir.) Sch.-Bip. (SI), *S. laniceps* Hand.-Mazz. (SL), *S. medusa* Maxim. (SM) plants and their medicinal materials.



**Fig. 2.** Anti-inflammatory effect of three herbal extracts on xylene-induced ear edema in mice. SI: *Saussurea involucrata*; SL: *Saussurea laniceps*; SM: *Saussurea medusa*. Values of edema shown are mean  $\pm$  S.E.M. ( $n=10$ ).  
\*  $p < 0.05$ ; \*\*  $p < 0.01$  vs. control group.




**Fig. 3.** UPLC chromatogram of a sample of SI extract (SI-E), a plasma sample of SI-treated rat (SI-R) and a plasma sample of SI-treated mouse (SI-M). Peak 1 to peak 3 were identified as chlorogenic acid, rutin and 1,5-dicaffeoylquinic acid, respectively.



**Fig. 4.** UPLC chromatogram of a sample of SL extract (SL-E), a plasma sample of SL-treated rat (SL-R) and a plasma sample of SL-treated mouse (SL-M).

Peak 1 to peak 7 were identified as umbelliferone 7-*O*- $\beta$ -D-glucoside, chlorogenic acid, scopoletin 7-*O*- $\beta$ -D-glucoside, syringoside, umbelliferone, scopoletin and 1,5-dicaffeoylquinic acid, respectively.



**Fig. 5.** UPLC chromatogram of a sample of SM extract (SM-E), a plasma sample of SM-treated rat (SM-R) and a plasma sample of SM-treated mouse (SM-M). Peak 1 to peak 3 were identified as chlorogenic acid, arctiin and arctigenin, respectively.